

Pest Alert



United States
Department of Agriculture

Forest Service

Animal and Plant
Health Inspection Service

NA-PR-01-99
Revised March 2010

Asian Longhorned Beetle (*Anoplophora glabripennis*): A New Introduction

The Asian longhorned beetle (ALB) has been discovered attacking trees in the United States. Tunneling by beetle larvae girdles tree stems and branches. Repeated attacks lead to dieback of the tree crown and, eventually, death of the tree. ALB probably travelled to the United States inside solid wood packing material from China. The beetle has been intercepted at ports and found in warehouses throughout the United States.

This beetle is a serious pest in China, where it kills hardwood trees in roadside plantings, shelterbelts, and plantations. In the United States the beetle prefers maple species (*Acer* spp.), including **boxelder, Norway, red, silver, and sugar maples**. Other preferred hosts are **birches, Ohio buckeye, elms, horsechestnut, and willows**. Occasional to rare hosts include **ashes, European mountain ash, London planetree, mimosa, and poplars**. A complete list of host trees in the United States has not been determined.

Currently, the only effective means to eliminate ALB is to remove infested trees and destroy them by chipping or burning. To prevent further spread of the insect, quarantines are established to avoid transporting infested

trees and branches from the area. Early detection of infestations and rapid treatment response are crucial to successful eradication of the beetle.

The ALB has one generation per year. Adult beetles are usually present from July to October, but can be found later in the fall if temperatures are warm. Adults usually stay on the trees from which they emerged or they may disperse short distances to a new host to feed and reproduce. Each female usually lays 35-90 eggs during her lifetime. Some are capable of laying more than that. The eggs hatch in 10-15 days. The larvae feed under the bark in the living tissue of the tree for a period of time and then bore deep into the wood where they pupate. The adults emerge from pupation sites by boring a tunnel in the wood and creating a round exit hole in the tree.

For more information about Asian longhorned beetle in the United States, visit these U.S. Department of Agriculture Web sites:

www.na.fs.fed.us/fhp/alb/

www.aphis.usda.gov/plant_health/plant_pest_info/asian_lhb/index.shtml

If you suspect an Asian longhorned beetle infestation, please collect an adult beetle in a jar, place the jar in the freezer, and immediately notify any of these officials or offices in your State:

State Department of Agriculture:

- State Plant Regulatory Official
- State Entomologist

U.S. Department of Agriculture:

- Animal and Plant Health Inspection Service, Plant Protection and Quarantine
- Forest Service

County Cooperative Extension Office

State Forester or Department of Natural Resources

Call 866-702-9938 toll free.

Asian Longhorned Beetle

WHAT TO LOOK FOR:



1. Adult beetles. Individuals are $\frac{3}{4}$ to $1\frac{1}{4}$ inches long, with jet black body and mottled white spots on the back. The long antennae are $1\frac{1}{2}$ to $2\frac{1}{2}$ times the body length with distinctive black and white bands on each segment. The feet have a bluish tinge.



2. Oval to round pits in the bark. These egg-laying sites or niches are chewed out by the female beetle, and a single egg is deposited in each niche.



3. Oozing sap. In the summer, sap may flow from egg niches, especially on maple trees, as the larvae feed inside the tree.



4. Accumulation of coarse sawdust around the base of infested trees, where branches meet the main stem, and where branches meet other branches. This sawdust is created by the beetle larvae as they bore into the main tree stem and branches.



5. Round holes, $\frac{3}{8}$ inch in diameter or larger, on the trunk and on branches. These exit holes are made by adult beetles as they emerge from the tree.

Photo Sources:

USDA Forest Service

USDA Animal and Plant Health Inspection Service

USDA is an equal opportunity provider and employer.



Published by:
USDA Forest Service
Northeastern Area
State and Private Forestry
Newtown Square, PA 19073
www.na.fs.fed.us



Federal Recycling Program
Printed on recycled paper.

ASIAN LONGHORNED BEETLE LOOK-ALIKES

(all insects shown approximately to scale, sizes given do not include antennae)

Asian longhorned beetle

(*Anoplophora glabripennis*)


$\frac{3}{4}$ – 1½ inches long

Shiny black, bright white spots

Long antennae, banded

black and white

Black scutellum



Whitespotted sawyer

(*Monochamus scutellatus*)


$\frac{3}{4}$ – 1 inch long

Dull or bronzy-black, may be

mottled with whitish patches

Long faintly banded antennae

White scutellum




Northeastern sawyer

(*Monochamus notatus*)

$\frac{3}{4}$ – 1½ inch long

Mottled light brown/white,

no distinct spots



Eyed click beetle

(*Alaus oculatus*)

1 - 1½ inch long

Black with white speckles;

Black "eye spots"




Brown prionid

(*Orthosoma brunneum*)

1 – 1½ inches long

Light chestnut brown, no spots




Broadnecked root borer

(*Prionus laticollis*)

1 – 1½ inches long

Black with no white markings

Solid black antennae



Longhorned Beetle

(*Graphisurus fasciatus*)

$\frac{1}{2}$ – $\frac{2}{3}$ inch long

Dark with gray/light brown mottling

Antennae banded white and black



Western conifer seed bug

(*Leptoglossus occidentalis*)

About $\frac{3}{4}$ inch long

Various shades of brown with a distinct geometric pattern on wings

Back legs flared-out at bottom



For more information about the Asian longhorned beetle, or to report a sighting, visit mass.gov/agr/alb.htm or call toll-free 1-866-702-9938



Flyer adapted from a design by the Maine Department of Agriculture

Photos: Asian longhorned beetle: Pennsylvania Dept. of Conservation and Natural Resources - Forestry Archive, Bugwood.org. Whitespotted sawyer, Northeastern sawyer, Broadnecked root borer: Michael Bohne, USDA Forest Service. Eyed click beetle: Tom Murray, bugguide.net. Brown prionid: Kristin Riolo, Priceless Memories, Bugwood.org. Graphisurus fasciatus: Jennifer Forman Orth, Massachusetts Dept. of Agricultural Resources. Western conifer seed bug: Giancarlo Dessì, Wikimedia Commons